

Attitude, Practice and Knowledge of Undergraduate Medical Students Towards Musculoskeletal Effects of Smoking in Saudi Arabia

Bandar Hetaimish (1)

Haneen Estanboli (2)

Asseil Bossei (2)

Ohood Shrouro (2)

Nizar Wali (2)

(1) MSc, FRCSC. Associate Professor of Orthopedic Surgery at University of Jeddah, Adjunct Professor of Orthopedic Surgery at Ibn Sina National College. Orthopedic Surgery Department, College of Medicine, University of Jeddah, Jeddah, Saudi Arabia

(2) Medical Intern at Ibn Sina National College for medical studies, Jeddah, Saudi Arabia

Corresponding author:

Bandar Hetaimish

Associate Professor of Orthopedic Surgery at University of Jeddah,

Adjunct Professor of Orthopedic Surgery, Ibn Sina National College,

P.O. Box 80327, post code 21589,

Jeddah, Saudi Arabia

Email: bmhetaimish@uj.edu.sa

Received: November 2019; Accepted: December 2019; Published: January 1, 2020.

Citation: Bandar Hetaimish, Haneen Estanboli, Asseil Bossei, Ohood Shrouro, Nizar Wali. Attitude, Practice and Knowledge of Undergraduate Medical Students Towards Musculoskeletal Effects of Smoking in Saudi Arabia. World Family Medicine. 2020; 18(1): 102-109. DOI: 10.5742/MEWFM.2020.93735

Abstract

Background: Smoking is a chief cause of inevitable death and disability. It is considered a risk factor for prospect fracture by decreasing bone density and has deleterious consequences on bone quality.

Objectives: The aims of this study were to determine the prevalence of tobacco smoking and assess the awareness of musculoskeletal effects of smoking among undergraduate medical and health science students at Makkah region, Kingdom of Saudi Arabia (KSA).

Method: A cross-sectional study was performed, using a questionnaire on a randomly selected number of students at Makkah region medical colleges in KSA. Data analysis was performed by a statistical team using SPSS program (version 16). The means and standard deviations of normally distributed variables were compared using paired t tests and for categorical variables, the X² test was used. The p-value of less than 0.05 was considered to be statistically significant.

Results: We had 370 participants, 56.6% female, and 31.3% male. Smoking prevalence among medical students reached 71.1% smokers, 39.3% smokers for more than 4 years while 60.7% smoked for less than 4 years and 28.9% did not smoke. 64.1% of medical students believe that smoking will affect bone health, while 13.2% did not believe and 22.7% did not know that there is any relationship.

Conclusion: This study identified that prevalence of smoking among undergraduate medical students at Makkah region is higher than previous reports in KSA. There is a crucial need to endorse multi-disciplinary health education events at different age groups to prevent adolescent students smoking, and to support smoking cessation programs.

Key words: Smoking, cigarette, bone health, fracture, medical students, Saudi Arabia

Introduction

Smoking has been recognized as the greatest significant reason of preventable disease and early mortality [1]. Though various adverse health effects of smoking happen later in the lifespan, smoking leads to complications in adolescent people as well [2]. Every day, roughly 4,800 adolescents smoke their first cigarette; of these, approximately 2000 will be changed to smokers [3]. Smoking-related diseases are attributed to smoking period (smoking years) and frequency (cigarettes/day). Furthermore, adult smokers started to smoke or had previously become habituated before 18 years of age [4]. Although many teenagers want to quit smoking, only a small number of them do [5]. Certainly, there has been an intense rise over the past decade in the total of college-age smokers [6]. Numerous researchers have stated that the prevalence of smoking rises from the fundamental to clinical years amongst medical college students, emphasizing the significance of directing anti-smoking activities to the fundamental years [7, 8]. For example, students who enter college as non-smokers are 40% less likely to start smoking if they live in a smoke-free campus [9]. With regard to Arab nations in particular, the World Health Organization has stated broadly distinctive prevalence rates of smoking amongst adolescent people: 18% in Kuwait, 43% in Yemen, 23% in Iraq, 25% in Kingdom of Saudi Arabia (KSA) and Jordan, 7% in Oman, 31% in Syrian Arab Republic and 53% in Lebanon [10]. Nevertheless, the pattern of smoking as well as the cessation rate, particularly amongst college students, is fundamentally unknown in many of these countries, including KSA. One study calculates the prevalence of active smoking amongst male medical students at King Saud University to be roughly 13% [11]. In 2009, Al-Turki et al discovered that the prevalence of smoking ranges from 2.4-52.3% among medical students in Central Saudi Arabia [11]. It is highlighted that nicotine has crucial side effects that may disturb most body systems, for example, the cardiovascular system, reproductive system, respiratory system, urinary system and also the musculoskeletal system [12]. Some of the side effects that can be caused by nicotine and carbon monoxide are decreasing the tissue oxygenation as well as micro-perfusion, and on the other hand, they also raise the rate of polycythemia and platelet aggregation [13]. Furthermore, the blood viscosity will increase while the total of oxyhemoglobin will be reduced due to carbon monoxide [13]. As a result, nicotine can affect the musculoskeletal system, predominantly bone healing. In 2016, Pearson, Clement, Edwards and Scammell showed that the risk of delayed or nonunion bone healing is 2.2 times greater in smokers. They explored that bone union time would take nearly 27.7 days longer in smokers than non-smokers [14]. Referring to clinical trials and demographic research which has been done throughout the countries, it disclosed that individuals who smoke have poor prognosis for fracture healing [15]. Furthermore, the negative impact of smoking on the bones is that it disturbs mineral density, lumbar disc degeneration and rate of hip fractures [15]. Smoking can lead to osteoporosis, spine and joint arthritis, devastate the cartilages and raise the risk of surgical infection [16].

There are a couple of research studies which have been made about the consequence of smoking on the musculoskeletal system, and bone healing process, in Saudi Arabia [16]. Fractures are a chief communal health concern, with estimates of over 3 million fractures yearly at a financial cost of \$25.3 billion by the year 2025 in the USA only [17]. Hip fractures provide an unreasonable burden on healthcare budgets and accompanying that it is not only an important disease but also causes an increased death rate. Research has revealed that smoking is related to an increase in fracture risk, predominantly at the hip, and existing smoking status is part of the World Health Organization Fracture Risk Assessment Tool (FRAX®) [18]. There are statistics from four big meta-analyses evaluating fracture risk in smokers. In their analysis, Law and Hackshaw [19] also anticipated hip fracture risk from 19 cohort and case-control studies with a sample size of 133,434 with 3,889 fractures. They found a significantly increased risk of hip fracture in female smokers, with increasing risk as persons aged. Lorentzon et al. studied 1,068 young men (average age 19 years) including 93 active smokers. Smokers had lower areal Bone Mineral Density (BMD) at the spine and hip than nonsmokers. After modification for age, height, weight, calcium intake, and physical activity, smokers had lower cortical bone size at the tibia and thinner cortices at mutually the radius and tibia than nonsmokers. [20] In addition, smokers had lower trabecular volumetric BMD at the tibia but no difference in cortical volumetric BMD.

The objectives of the current study were to determine the prevalence of tobacco smoking and to assess the awareness of musculoskeletal effects of smoking among undergraduate medical and health science students at Makkah region, Kingdom of Saudi Arabia (KSA).

Methods

Design: A descriptive/ analytic cross-sectional study was performed, using a questionnaire on a sample size of 370 male and female students who were randomly selected at Western region medical colleges of Saudi Arabia.

Analysis: Data analysis was performed by statistical team using SPSS program (version 16). The means and standard deviations of normally distributed variables were compared using paired t tests and for categorical variables, the X² test was used. The p-value of less than 0.05 was considered to be a statistically significant.

Participants: Undergraduate medical students enrolled at Medical colleges in Makkah region who voluntarily responded to participate in the online survey.

Survey Instrument: After obtaining ethical approval from research ethic board committee at our institution, randomly selected consenting participants were asked to fill out a 25 items self-structured online questionnaire. It was first directed to 12 students of our college and pilot tested. Appropriate adjustments were then made before confirming it for the study. The questionnaire contained items to look for information regarding demography,

prevalence, and smoking pattern. Students were assured about the anonymity of their answers. Since knowledge of musculoskeletal effects of smoking evolves as we grow the survey included questions about respondents' sociodemographic, clinical information, education, history and pattern of smoking, and students' knowledge and beliefs of musculoskeletal effects of smoking, bone fractures, bone healing and physical activities. The questionnaire included primarily close-ended questions. Some of the questions allowed more than 1 answer. The questionnaire was settled after a comprehensive appraisal of the related

articles and consultation amongst the research team. It was face-validated through discussion with professional collaborators in the field and was moreover objectively validated for comprehensibility.

Implications of results:

Results will be developed into educational awareness planning and interventions for incoming undergraduate students.

Results

Table 1: Prevalence of smoking in medical school

Do you smoke?	Medical school	Mean	N	Std. Deviation
Yes	ISNC	1.33	53	.474
	UQU	1.00	1	.
	KAU	1.55	20	.510
	Farabi	1.50	2	.707
	Taif	1.00	3	.000
	Fakeeh	1.50	2	.707
	BMC	1.40	5	.548
	KSAU-HS	1.50	4	.577
	Other	1.06	18	.236
	Total	1.33	108	.471
No	ISNC	1.74	102	.443
	UQU	1.53	15	.516
	KAU	1.86	59	.345
	Farabi	1.60	5	.548
	Taif	1.81	16	.403
	Fakeeh	1.00	1	.
	BMC	1.88	8	.354
	KSAU-HS	1.50	4	.577
	Other	1.44	52	.502
	Total	1.69	262	.461
Total	ISNC	1.60	154	.492
	UQU	1.50	16	.516
	KAU	1.78	79	.414
	Farabi	1.57	7	.535
	Taif	1.68	19	.478
	Fakeeh	1.33	3	.577
	BMC	1.69	13	.480
	KSAU-HS	1.50	8	.535
	Other	1.34	70	.478
	Total	1.59	369	.493

ISNC: Ibn Sina National College for Medical Studies, UQU: Umm Al-Qura University, KAU: King Abdulaziz University, Farabi: Al-Farabi college, Taif: Taif university, Fakeeh: Fakeeh College for Medical Sciences, BMC: Batterjee Medical College, KSAU-NG: King Saud Bin Abdulaziz University for Health Science

Table 2: Knowledge of smoking effect on bone and general health

Have you ever experienced bone pain ?	Which bone had you fractured ?	Have you ever had bone fracture ?	Mean	N	Std. Deviation	
Yes	Forearm	Yes	1.00	1	.	
		No	1.00	1	.	
		Total	1.25	8	.463	
	Fingers	Yes	1.00	1	.	
		No	1.48	67	.503	
		Total	1.45	71	.501	
	Total	Yes	1.00	2	.000	
		No	1.47	69	.503	
		Total	1.43	79	.498	
No	Foot	non	1.63	290	.483	
		Total	1.63	290	.483	
	Total	non	1.63	290	.483	
		Total	1.63	290	.483	
Total	Forearm	Yes	1.00	1	.	
		No	1.00	1	.	
		Total	1.25	8	.463	
	Fingers	Yes	1.00	1	.	
		No	1.48	67	.503	
		Total	1.45	71	.501	
	Foot	non	1.63	290	.483	
		Total	1.63	290	.483	
	Total	Yes	Yes	1.00	2	.000
			No	1.47	68	.503
		Total	non	1.63	290	.483
			Total	1.59	369	.493

Table 3: Relation of smoker to general exercising and health activity

Do you smoke?	Do you perform exercise/ physical activities ?	Mean	N	Std. Deviation
Yes	1-2 days	1.37	28	.492
	3-4 days	1.32	25	.476
	5-7 days	1.15	13	.376
	0	1.36	42	.485
	Total	1.33	108	.471
No	1-2 days	1.69	55	.466
	3-4 days	1.68	68	.471
	5-7 days	1.69	35	.471
	0	1.71	104	.455
	Total	1.69	262	.461

Summary of results

The prevalence of smoking in our sample was 71.1% and 28.9% are non-smoker. 39.3% of smokers had smoked for a period of more than 4 years while 60.7% smoked for less than 4 years.

Our results showed that a bulk of students who smoke represents 49.5% and they reported that they smoke cigarettes and 25.3% smoke shisha while 18.7% smoke Dokha (Arabian tobacco product, consisting of dried and finely shredded tobacco flakes mixed with herbs and spices). Nevertheless, 10.3% smoked more than 20 cigarettes per day, 41.1% of smokers smoked 5 cigarettes per day, while the rest, 48.6%, smoked 10 – 20 cigarettes per day.

We found that the majority, 39.3%, of smokers don't want to quit. On the other hand, we found that 36.4% plan to quit smoking, and 24.3% plan to quit after finishing medical school. There are numerous potential explanations for the extraordinary prevalence, including high pressure of medical specialty. Overall, smoking and physical activity seems to be negatively associated, but such simplifications must be made with caution as there may be many causes.

We found in our study that 60.5% perform exercise, 40.2% of them spend 30 minutes, 25% spend 60 minutes, 19.6% spend 10 minutes and 15.2% spend more than 60 minutes, while 39.5% don't exercise. On the other hand, 42% from those who performed exercise do physical activities 3-4 days a week as a part of their work and 36.6% do 1-2 days a week while 21.4% do it 5-7 days.

On other hand, 68.1% of students know that smoking increases post-surgical wound healing complications risk whereas 31.9% did not know that risk. The main bulk of undergraduate medical students 51.9% do not know that smoking destroys cartilage while 48.1% knew that fact. 54.6% of students know that smoking delays healing of tendons repair while 45.4% of students are not familiar with that fact.

Discussion

Our study offered insight about the prevalence and attitudes in respect to Musculoskeletal effects of smoking in Saudi Arabian medical students from different specialties in Jeddah, Saudi Arabia. The prevalence of smoking in our sample was 71.1% and 28.9% were non-smokers. 39.3% of smokers smoked for a period of more than 4 years while 60.7% smokers for less than 4 years. The prevalence is greater than the prevalence of 27.8% stated in 2014 amongst dental students at King Saud University, KSA [21]. In addition, that number is higher than the 24.8% prevalence amongst undergraduate medical students in the western region of KSA [22] and the 17.6% amongst undergraduate medical students at King Fahad Medical City in Riyadh, KSA [23]. This outcome is also greater than the results of an article conducted between students at a Malaysian college, which stated that the prevalence

was 29% [24]. Moreover, our result is higher compared to a study of smoking amongst Jordan University medical students that revealed a total prevalence of 50.2% [25]. Our results showed that the bulk of students who smoke represents 49.5% and they reported that they smoked cigarettes and 25.3% smoked shisha while 18.7% smoked Dokha. Nevertheless, 10.3% smoked more than 20 cigarettes per day, 41.1% of smokers smoked 5 cigarettes per day, while the rest, 48.6%, smoked 10 – 20 cigarettes per day. This outcome varies from the results of the national analysis of the general public in the kingdom that revealed that commencement of smoking was more common at the age of 19 years [26]. At such an age, students are likely to be in university. We found that the majority (39.3%) of smokers don't want to quit. On the other hand, we found that 36.4% have a plan to quit smoking, and 24.3% have a plan to quit after finishing medical school. There are numerous potential explanations for the extraordinary prevalence, including high pressure of medical specialty. Overall, smoking and physical activity seems to be negatively associated, but such simplifications must be made with caution for many causes. Though a bulk of studies advocate a reverse relationship between physical activity and smoking, this relationship seemed to be more attenuated in youths, and multifaceted relations may occur for other people subgroups [27]. We found in our study that 60.5% perform exercise, 40.2% of them spend 30 minutes, 25% spend 60 minutes, 19.6% spend 10 minutes and 15.2% spend more than 60 minutes, while 39.5% do not perform exercise. On the other hand, 42% from those who performed exercise do physical activities 3-4 days a week as a part of their work and 36.6% do it 1-2 days while 21.4% do it 5-7 days. There are hypothetical primary and secondary special effects of smoking on musculoskeletal health and risk of fracture. Primary toxic consequences of smoking on bone may be associated with nicotine special effects [28, 29] or perhaps to toxic compounds in tobacco products like cadmium [30]. Smoking has direct special effects on osteogenesis involving change in the RANK–RANKL–OPG system [31,32], collagen metabolism [33], and bone angiogenesis [34]. Secondary special effects of smoking on bone might result from decreased calcium absorption from intestine [35], sex hormone dysregulation in production [36], cortical and gonadal hormones metabolism alterations [37–39], calcitropic hormones [40] like 25-hydroxy vitamin D [36, 41] plus parathyroid hormone [36]. These consequences may explain the commonly observed decrease in indications of bone formation, such as osteocalcin, among smokers [41, 42]. Smoking also has indirect influence on bone density and fractures risk through reductions in body weight. Body weight tends to be less for smokers than non-smoking individuals, and this weight differentiation may lead to lower bone density and increased fracture risk [43, 44]. Ultimately, smokers might be less physically active, which may decrease bone density [45] and increase risk of fracture [46]. Certain elements such as high BMI [47], and high calcium consumption [48] have been described to attenuate the smoking relationship with bone. Mosely and Finseth, found that smoking had a harmful effect on hand wounds healing. The adversarial effect of smoking

on fracture healing has been the base of much clinical research, and there is a strong relationship with cardiac and pulmonary diseases [49–54]. On other hand, 68.1% of students know that smoking increases post-surgical wound healing complications risk whereas 31.9% do not know that risk. The main bulk of undergraduate medical students 51.9% do not know that smoking destroys cartilage while 48.1% know that fact. 54.6% of students know that smoking delays healing of tendons repair while 45.4% do not. Research results have suggested that nicotine and related substances in cigarettes can also impair the regeneration of wound healing and soft tissues after fracture, thus decreasing the quality of postoperative consequences and delaying wound healing [55–57]. Our study shows that 59.5% of students know that smoking is a risk factor for osteoporosis while 50.5% do not know. 64.1% of students believe that smoking will affect bone health while 13.2% do not believe that and 22.7% do not know that there is a relationship. Regarding bone fractures, 78.6% did not experience bone fracture while 21.4% did and the commonest fracture was in the arm with 30.4%. There are over a million bone fractures each year in the United Kingdom, and 5–10% are stated not to heal adequately. Thus it is critical that the orthopedic surgeon is aware of the risk factors that could potentially impair bone healing, in order to avoid them whenever possible when managing fractures. There are several theories as to how smoking can influence the healing process of bone fractures and incorporate a reduced blood supply to the injury site, high levels of reactive oxygen species (ROSs) in circulation, low levels of vitamins and antioxidants and the attenuating consequence nicotine has on synthase of endothelial nitric oxide. Bone health is affected by cigarette smoke, and is well known to augment osteoporosis and osteonecrosis of femur in both genders. [58] Whatever the mechanism, this information suggests that the fractures risk is higher for smokers and those who have a smoking history than it is for individuals of the same demographic and BMD who don't or did not smoke ever in their life. Nevertheless, a big number of stronger independent fracture risk factors have been recognized in previous reports. These include fracture history, prolonged use of corticosteroids, significant family history of fracture, secondary osteoporosis, and perhaps the biochemical indicators of bone turnover. These risk factors can be readily used for measuring risk of fracture in the community and their relationships to smoking will need to be determined [59].

Limitations

The study was built on self-reported information hence elicit bias cannot be ruled out. Also, some undergraduate medical students may not disclose their smoking status, nevertheless they were told that their data would be kept confidential.

Conclusion

This study has shown that the prevalence of smoking among health sciences students at Makkah region is higher than the prevalence of smoking reported by other studies in KSA. As the number of smokers globally continues to increase, we must assume an increased disease burden attributable to smoking, including an increased number of osteoporotic fractures. There is a demand to encourage multi-disciplinary health education activities at different age groups to prevent young medical students from smoking, and to assist smoking cessation programs.

Recommendation

1. Campaigns should be developed to raise public awareness of the benefits of cessation and available therapeutic options, including addressing misconceptions about the safety and effectiveness of treatments.
2. Educate the students and provide their parents with the necessary knowledge to educate their children on the danger they face.
3. Focus on the application of the basic principles, the most important of which are:
 - a. Implementing and activating the smoking prevention law within public and public institutions and buildings, and allocating limited sections for smokers in restaurants and cafes.
 - b. Increase the health warning on smoking boxes to include at least one third of the box space and add the warning image.
4. Work to provide specialist smoking cessation services within free health centers.
5. Pay attention to the high prevalence of smoking Shisha and work on preventing its promotion, and in particular promote the claim that it is less harmful than cigarette.

Conflict of Interest

Authors have no conflict of interest.

Acknowledgement

We would like to express our deep gratitude to Ms. Yumna Abdulmalek Bokhari, MBBS student at Ibn Sina National College for Medical Studies who assisted the research team in collecting and entering the research. We extend gratitude to all the participants in this study.

Financial support and sponsorship

There was no funding obtained for this research project by all of the authors.

References

1. WHO. Combating the tobacco epidemic, World Health Organization (Ed.), Making a difference in people's lives. World Health Organization, Geneva (1999)
2. Morbidity and mortality weekly report. Symptoms of substance dependence associated with use of cigarettes, alcohol, and illicit drugs. United States; 1995.
3. American Lung Association, Adolescent smoking statistics.
4. Center for Health Promotion. Youth and tobacco: preventing tobacco use among young people: a report of the Surgeon General Centers for Disease Control and Prevention. Atlanta, Georgia (1994)
5. L. Lamkin, T.P. Houston. Nicotine dependency and adolescents: preventing and treating. *Prim Care*, 25 (1998), pp. 123-135
6. T.P. Houston, L.J. Kolbe, M.P. Eriksen. Tobacco-use cessation in the 90s-not "adults only" anymore. *Prev Med*, 27 (1998), pp. A1-A2
7. D. Aslan, et al. Prevalence and determinant of adolescent smoking in Ankara, Turkey. *Turk J Cancer*, 36 (2) (2006), pp. 49-56
8. S. Yegenoglu, et al. What is behind smoking among pharmacy students: a quantitative and qualitative study from Turkey? *Subst Use Misuses*, 41 (2006), pp. 405-414
9. Faculty Senate Special report of the health council concerning a tobacco free campus (2003)
10. Middle East and North Africa. 11-Economics of tobacco for the Middle East and North Africa (MNA) region.
11. Yousef A. Al-Turki. Smoking habits among medical students in Central Saudi Arabia. *Saudi Med J*, 27 (5) (2006), pp. 700-703
12. Mishra, A., Chaturvedi, P., Datta, S., Sinukumar, S., Joshi, P., & Garg, A. (2015). Harmful effects of nicotine. *Indian Journal of Medical and Paediatric Oncology: Official Journal of Indian Society of Medical & Paediatric Oncology*, 36(1), 24-31.
13. Lee, John J. MD, MS; Patel, Rakesh MD; Biermann, J. Sybil MD; Dougherty, Paul J. MD. (2013). The musculoskeletal effects of cigarette smoking. *The Journal of Bone and Joint Surgery*, 95(9), 850-59. doi: 10.2106/JBJS.L.00375
14. Davies, T, M., Wluka, A, E., Forbes, A., Wang, Y., English, D, R., Giles, G, G., and Cicuttini, F. (2009). Smoking is associated with increased cartilage loss and persistence of bone marrow lesions over 2 years in community-based individuals. *Rheumatology*, 48(10), 1227-1231. DOI: 10.1093/rheumatology/kep211
15. Sloan, A., Hussain, I., Maqsood, M., Ermin, O., and El-Sheemy, M. (2010). The effects of smoking on fracture healing. *Surge*, 8 (2), 111-16. DOI: 10.1016/j.surge.2009.10.014
16. Bassiony, M., M. (2009). Smoking in Saudi Arabia. *Saudi Med J*, 30(7), 876-881.
17. Burge R, Dawson-Hughes B, Solomon DH, Wong JB, King A, Tosteson A. Incidence and economic burden of osteoporosis related fractures in the United States, 2005-2025. *J Bone Miner Res Off J Am Soc Bone Miner Res*. 2007;22(3):465-75.
18. Kanis JA, Johnell O, Oden A, Johansson H, De Laet C, Eisman JA, et al. Smoking and fracture risk: a meta-analysis. *Osteoporosis Int J Established Result Cooperation Between Eur Found Osteoporosis Natl Osteoporosis Found USA*. 2005;16(2):155-62.
19. Law MR, Hackshaw AK. A meta-analysis of cigarette smoking, bone mineral density and risk of hip fracture: recognition of a major effect. *BMJ*. 1997;315(7112):841-6
20. Lorentzon M, Mellstrom D, Haug E, Ohlsson C. Smoking is associated with lower bone mineral density and reduced cortical thickness in young men. *J Clin Endocrinol Metab*. 2007;92(2):497-503.
21. A.S. AlSwuaillem, M.K. AlShehri, S. Al-Sadhan. Smoking among dental students at King Saud University: consumption patterns and risk factors. *Saudi Dent J* (2014), pp. 88-95, 10.1016/j.sdentj
22. Siraj O. Wali. Smoking habits among medical students in Western Saudi Arabia. *Saudi Med J*, 32 (8) (2011), pp. 843-848
23. A.F. Al Kaabba, A.A. Saeed, A.M. Abdalla, H.A. Hassan, A.A. Mustafa. Prevalence and associated factors of cigarette smoking among medical students at King Fahad Medical City in Riyadh of Saudi Arabia. *J Fam Community Med* (2011), pp. 8-12, 10.4103/1319-1683.78631
24. R.A. Al-Naggar, S.A. Al-Dubai, T.H. Al-Naggar, R. Chen, K. Al Jashamy. Prevalence and of smoking and associated factors among Malaysian University students. *Asian Pac J Cancer Prev*, 12 (3) (2011), pp. 619-624
25. Linda G. Haddad, Malakeh Z. Malak. Smoking habits and attitudes towards smoking among university students in Jordan. *Int J Nurs Stud*, 39 (2002), pp. 793-802
26. M. Moradi-Lakeh, C. ElBcheraoui, M. Tuffaha, F. Daoud, M. Al Saeedi, M. Basulaiman, Z.A. Memish, M.A. AlMazroa, A.A. AlRabeeah, A.H. Mokdad. Tobacco consumption in the Kingdom of Saudi Arabia, 2013: findings from a national survey. *BMC Public Health*, 15 (2015), p. 611, 10.1186/s12889-015-1902-3
27. Kaczynski, A. (2008). Smoking and Physical Activity: A Systematic Review. *American Journal of Health Behavior*, 32(1). doi:10.5993/ajhb.32.1.9
28. Riebel GD, Boden SD, Whitesides TE, Hutton WC. The effect of nicotine on incorporation of cancellous bone graft in an animal model. *Spine*. 1995;20(20):2198-202.
29. Fang MA, Frost PJ, Iida-Klein A, Hahn TJ. Effects of nicotine on cellular function in UMR 106-01 osteoblast-like cells. *Bone*. 1991;12(4):283-6.
30. Bhattacharyya MH, Whelton BD, Stern PH, Peterson DP. Cadmium accelerates bone loss in ovariectomized mice and fetal rat limb bones in culture. *Proc Natl Acad Sci U S A*. 1988;85:8761-5.
31. Tang TH, Fitzsimmons TR, Bartold PM. Effect of smoking on concentrations of receptor activator of nuclear factor kappa B ligand and osteoprotegerin in human gingival crevicular fluid. *J Clin Periodontol*. 2009;36(9):713-8.
32. Lappin DF, Sherrabeh S, Jenkins WM, Macpherson LM. Effect of smoking on serum RANKL and OPG in sex, age and clinically matched supportive-therapy periodontitis patients. *J Clin Periodontol*. 2007;34(4):271-7.

33. Sorensen LT, Toft BG, Rygaard J, Ladelund S, Paddon M, James T, Taylor R, et al. Effect of smoking, smoking cessation, and nicotine patch on wound dimension, vitamin C, and systemic markers of collagen metabolism. *Surgery*. 2010;148(5):982–90.
34. Ma L, Zheng LW, Sham MH, Cheung LK. Uncoupled angiogenesis and osteogenesis in nicotine-compromised bone healing. *J Bone Miner Res*. 2010;25(6):1305–13.
35. Krall EA, Dawson-Hughes B. Smoking increases bone loss and decreases intestinal calcium absorption. *J Bone Miner Res*. 1999;14(2):215–20.
36. Supervia A, Nogues X, Enjuanes A, Vila J, Mellibovsky L, Serrano S, Aubia J, et al. Effect of smoking and smoking cessation on bone mass, bone remodeling, vitamin D, PTH and sex hormones. *J Musculoskelet Neuronal Interact*. 2006;6(3):234–41.
37. Baron JA, Comi RJ, Cryns V, Brinck-Johnsen T, Mercer NG. The effect of cigarette smoking on adrenal cortical hormones. *J Pharmacol Exp Ther*. 1995;272(1):151–5.
38. Khaw KT, Tazuke S, Barrett-Connor E. Cigarette smoking and levels of adrenal androgens in postmenopausal women. *N Engl J Med*. 1988;318(26):1705–9.
39. Michnovicz JJ, Hershcopf RJ, Naganuma H, Bradlow HL, Fishman J. Increased 2-hydroxylation of estradiol as a possible mechanism for the anti-estrogenic effect of cigarette smoking. *N Engl J Med*. 1986;315(21):1305–9.
40. Yoon V, Maalouf NM, Sakhaee K. The effects of smoking on bone metabolism. *Osteoporos Int*. 2012;23(8):2081–92.
41. Brot C, Jorgensen NR, Sorensen OH. The influence of smoking on vitamin D status and calcium metabolism. *Eur J Clin Nutr*. 1999;53(12):920–6.
42. Bjarnason NH, Christiansen C. The influence of thinness and smoking on bone loss and response to hormone replacement therapy in early postmenopausal women. *J Clin Endocrinol Metab*. 2000;85(2):590–6.
43. Cummings SR, Nevitt MC, Browner WS, Stone K, Fox KM, Ensrud KE, Cauley J, et al. Risk factors for hip fracture in white women. Study of Osteoporotic Fractures Research Group. *N Engl J Med*. 1995;332(12):767–73.
44. Kiel DP, Felson DT, Anderson JJ, Wilson PW, Moskowitz MA. Hip fracture and the use of estrogens in postmenopausal women. The Framingham Study. *N Engl J Med*. 1987;317(19):1169–74.
45. Bolam KA, van Uffelen JG, Taaffe DR. The effect of physical exercise on bone density in middle-aged and older men: a systematic review. *Osteoporos Int*. 2013;24(11):2749–62.
46. Feskanich D, Willett W, Colditz G. Walking and leisure-time activity and risk of hip fracture in postmenopausal women. *JAMA*. 2002;288(18):2300–6.
47. Baheiraei A, Pocock NA, Eisman JA, Nguyen ND, Nguyen TV. Bone mineral density, body mass index and cigarette smoking among Iranian women: implications for prevention. *BMC Musculoskelet Disord*. 2005;6:34.
48. Ilich JZ, Brownbill RA, Tamborini L, Crncevic-Orlic Z. To drink or not to drink: how are alcohol, caffeine and past smoking related to bone mineral density in elderly women? *J Am Coll Nutr*. 2002;21(6):536–44.
49. Rees TD, Liverett DM, Guy CL. The effect of cigarette smoking on skin-flap survival in the face lift patient. *Plast Reconstr Surg* 1984 Jun;73(6):911–5.
50. Riefkohl R, Wolfe JA, Cox EB, McCarty Jr KS. Association between cutaneous occlusive vascular disease, cigarette smoking, and skin slough after rhytidectomy. *Plast Reconstr Surg* 1986 Apr;77(4):592–5.
51. Kroll SS. Necrosis of abdominoplasty and other secondary flaps after TRAM flap breast reconstruction. *Plast Reconstr Surg* 1994 Oct;94(5):637–43.
52. Bailey MH, Smith JW, Casas L, Johnson P, Serra E, de la Fuente R, et al. Immediate breast reconstruction: reducing the risks. *Plast Reconstr Surg* 1989 May;83(5):845–51.
53. Hartrampf Jr CR, Bennett GK. Autogenous tissue reconstruction in the mastectomy patient. A critical review of 300 patients. *Ann Surg* 1987 May;205(5):508–19.
54. Mosely LH, Finseth F. Cigarette smoking: impairment of digital blood flow and wound healing in the hand. *Hand* 1977 Jun;9(2):97–101.
55. Pitts KR, Yoon Y, Krueger EW, McNiven MA. The dynamin-like protein DLP1 is essential for normal distribution and morphology of the endoplasmic reticulum and mitochondria in mammalian cells. *Mol Biol Cell* 1999 Dec;10(12):4403–17.
56. Netscher DT, Clamon J. Smoking: adverse effects on outcomes for plastic surgical patients. *Plast Surg Nurs* 1994 Winter;14(4):205–10.
57. Nolan J, Jenkins RA, Kurihara K, Schultz RC. The acute effects of cigarette smoke exposure on experimental skin flaps. *Plast Reconstr Surg* 1985 Apr;75(4):544–51.
58. Delmas PD, Eastell R, Garnero P, Seibel MJ, Stepan J (2000). The use of biochemical markers of bone turnover in osteoporosis. *Osteoporos Int [Suppl 6] 11:S2-S17*
59. Kanis, J. A., Johnell, O., Oden, A., Johansson, H., De Laet, C., Eisman, J. A., Tenenhouse, A. (2004). Smoking and fracture risk: a meta-analysis. *Osteoporosis International*, 16(2), 155–162. doi:10.1007/s00198-004-1640-3